


# FIBER CONSUMPTION REDUCES THE RISK OF COLORECTAL CANCER

by Charles B. Simone, M.D., Nicole L. Simone, and Charles B. Simone, II



Gastrointestinal cancers are the second leading cause of death among all cancer patients. The death rate for cancer of the colon and rectum has remained virtually the same since 1930, which means there has been essentially no progress in the treatment of these cancers. The new colorectal cancer cases for 2000 were approximately 130,200. The estimated number of deaths from colorectal cancer is 56,600. A person has a one-in-18 chance of developing colorectal cancer<sup>1</sup> over his or her lifetime.

Major differences in death rates from colorectal cancer occur in different parts of the world, and epidemiological studies show that dietary factors account for the different incidence rates.<sup>2</sup> The more industrialized a country, the higher the rate of colorectal cancer

because the people generally eat less fiber and more animal fat. The highest colorectal cancer rates are found in Western Europe and English-speaking countries. The lowest rates are found in Africa and Asia, but that is changing rapidly for Asians who have adopted a Westernized diet.

In countries with a high incidence of colorectal cancer, most of the cancers are located in the left colon and rectum, whereas in countries with a low incidence, most of the cancers are in the right colon. Carcinogens become progressively more concentrated at the end of the gastrointestinal tract (left colon and rectum). Dr. Denis Burkitt shares the following analogy for this distribution of cancer: "While a man proceeds down a path carrying a leaky pot of water containing a tablet

of dye that is gradually dissolving, the water becomes more deeply colored because the volume will be progressively reduced and the dye more concentrated, and more dye will be progressively dissolved."

### ROLE OF FIBER

Fiber is a complex carbohydrate consisting of a polysaccharide and a lignin substance that provides the structure of a plant cell. It is undigested residue that reaches the end of the small intestine. The three groups of dietary fiber types are vegetable fibers, which are highly fermentable and have a low undigested content; bran, which is less fermentable; and purified fibers, such as cellulose, which are much less fermentable and have a high undigested content.

Dietary fiber acts as a "glue" for certain chemicals. For instance, unconjugated bile acids, which the body produces, can be absorbed to fiber in the colon and passed out in the stool without intestinal bacteria forming carcinogens from those bile acids. In addition, some fiber binds to cholesterol, lipids, nitrogen, and certain minerals, and eliminates them in the stool. This action lowers the blood concentration of cholesterol and certain other lipids.

Drs. Higginson and Oettle were the first to report in 1960 that dietary fiber consumption was associated with a low risk of developing colon cancer.<sup>3</sup> They noted that the Bantu tribal people in South Africa had a low incidence of colorectal cancer. They also excreted large piles of feces that were related to the large amount of dietary fiber they ate. Dr. Denis Burkitt continued the research and concluded that the high-fiber diet resulted in a rapid transit time for solid material to pass through the gastrointestinal tract and also increased the amount of stool. These two variables are associated with a decreased incidence of colorectal cancer.<sup>4</sup>

Diet in rural Africa and in other similar locations provides about 25 grams of crude fiber daily, whereas Western diets provide only 8 to 15 grams of fiber daily. With a more rapid transit time, bile acids and other carcinogens produced by anaerobic bacteria move out of the gastrointestinal tract more quickly. Furthermore, since the volume of feces is increased, carcinogens that are produced pass through the gut more diluted. Hence, if more dietary fiber is eaten, carcinogens pass out of the gut more quickly and there are fewer carcinogens per square inch.

### EVALUATING THE DATA

In 1982, the National Academy of Sciences found that, according to strict epidemiological criteria, there was "no conclusive evidence to indicate that dietary fiber exerts a protective effect against colorectal cancer in humans." Nevertheless, the U.S. National Academy of Sciences did issue dietary guidelines because the data were "highly suggestive that reduced fat consumption and increased consumption of cereals, fruit, and vegetables represent the current state of knowledge and form the basis of a diet that

is unlikely to do harm and may have the potential for reducing cancer rates in North America."

Because the evidence from epidemiological and laboratory studies was sufficiently consistent that high-fiber, low-fat diets could lower cancer risk, other U.S. agencies, organizations, and other governments issued interim dietary guidelines in the mid 1980s. These included the United States National Cancer Institute, National Institutes of Health, United States Department of Agriculture, Department of Health and Human Services, American Cancer Society, Australia, Canada, the Joint European Organization for Cooperation in Cancer Prevention, Norway, Sweden, and Japan. They all independently agreed that to reduce cancer risk, people should increase their consumption of green, yellow, and cruciferous vegetables, citrus fruits, and whole-grain cereal products; and reduce their intake of fats to about 30 percent.

In 1984, the United States National Cancer Institute recommended an intake of 25-35 grams of fiber daily to decrease the risk of cancer. However, the American public consumes only about 8-15 grams of fiber per day.

During the last 25 years, thousands of *in vitro* and animal studies have been published demonstrating that fiber can decrease the risk of colorectal cancer. These papers have not been included in this review. Since 1980, hundreds of published papers demonstrate that high fiber intake can reduce the incidence of colorectal cancer in humans. Some have been included in this review.<sup>5-108</sup>

### REPORTS ON FIBER FROM THE NCI

Reports from the United States National Cancer Institute are consistent that dietary fiber can decrease the risk of colorectal cancer:

- "This evaluation clearly suggests a relationship between colon cancer and a diet low in fiber." From 40 epidemiological studies in 55 reports (Greenwald P, et al: Dietary fiber in the reduction of colon cancer risk. *J Am Diet Assoc* 87[9]:1178-1188, 1987).
- "The analysis of these studies gives support for a protective effect [against colorectal cancer] associated with fiber-rich diets." From 23 case-control studies, 7 international correlation studies, 8 within-country correlation studies, 2 cohort studies, and 3 time-trend studies (Trock, Lanza, Greenwald: Dietary fiber, vegetables, and colon cancer: critical review and meta-analysis of epidemiological studies. *JNCI* 82:650-661, 1990).
- "Based on current knowledge, recommended nutrition guidelines for reducing the risk of colon cancer include decreased fat consumption, adequate amounts of fruits, vegetables, and calcium, and avoidance of overweight" (Shike, Winawer, Greenwald, et al: Primary prevention of colorectal cancer. *Bull WHO* 68:377-385, 1990).
- "Both prospective and retrospective studies suggest that vegetable and fruit intake may reduce the risk of cancers...including cancer of the colon and rectum" (Ziegler RG: Vegetables, fruits, and carotenoids and the risk of cancer. Environmental Epidemiology Branch, National Cancer Institute, Bethesda, MD 20892. *Am J Clin Nutr* 53[1 Suppl]:251S-259S, January 1991).



Various organizations and governments around the world have issued consensus statements that high fiber consumption can reduce the risk of colorectal cancer.

- 1999 World Health Organization: "The consumption of foods rich in polysaccharides (e.g., dietary fiber or non-starch polysaccharides) is associated with a decreased risk of colorectal adenoma and colorectal cancer" (*Eur J Canc Prev* 3:57-62, 1999).

Recommendation: Vegetables and whole-grain cereals should be consumed in high amounts and should be a major component of the diet.

- 1999 Colon Cancer Prevention Program Project: "13.5 grams of wheat bran per day decreases the recurrence rate of adenomatous colon polyps" (*Am J Med* 106[1A]:43S-45S, 1999).

- 1999 The Seven Countries Study Conclusion (Croatia, Finland, Greece, Italy, Japan, Netherlands, Serbia, U.S.): "High fiber intake was strongly associated with low colorectal mortality. An increase of 10 grams in the daily intake of fiber was associated with a 33% lower risk of 25-year colorectal cancer mortality" (*Int J Cancer* 84:174-179, 1999).

Recommendation: Increase the daily intake of fiber by 10 grams.

- 1998 European Cancer Prevention Consensus Panel: "A diet rich in high-fiber cereal is associated with a reduced risk of colorectal cancer" (*Eur J Canc Prev* 7[suppl 2]:S1-S3, 1998).

- 1997 American Dietetic Association Position: "Results of all studies provide substantive evidence that intake of fiber-rich foods is inversely related to risks of both colon and rectal cancers. It is estimated that the risk of colorectal cancer in the U.S. population could be reduced by about 31 percent if fiber intake from food sources were increased by an average of about 13 grams per day" (*J Am Diet Assoc* 97[10]:1157-1159, 1997).

Recommendation: Promote food intake patterns consistent with the Food Guide Pyramid. This recommends a wide variety of plant foods to achieve adequate fiber intakes in healthy children and adults. Include at least 2 to 3 servings of whole grains as part of the daily 6 to 11 servings of grains, 2 to 4 servings of fruits, and 3 to 5 servings of vegetables daily, and legumes at least once or twice a week.

- 1995 Australia: "Reduction in the incidence in large adenomas was observed when a low-fat diet was combined with high-fiber wheat bran supplementation of 25 grams per day" (*JNCI* 87:1760-1766, 1995).

Recommendation: 25 grams of fiber daily.

- 1994 United Nations Food and Agriculture Organization: "High fiber intake consisting of vegetables and cereals was protective against colorectal cancer" (*Eur J Canc Prev* 7[suppl 2]:S11-S17, 1998).

#### RISK/BENEFIT OF

##### DAILY CONSUMPTION OF FIBER

According to the National Academy of Science, over 40% of North Americans are likely to develop cancer and at least half of them will die from it. The cancer incidence worldwide is increasing. The majority of health budgets will be spent on treating cancer in most developing countries.

Cancer is largely preventable. Fewer than 5% of cancer cases are linked to genetics.

Overwhelming evidence supports the statement that "the consumption of fiber may reduce the risk of colorectal cancer." In fact, based on the volume, credibility, and reliability of the scientific facts, we are convinced that fiber *can*, not *may*, but *can* reduce the risk of colorectal cancer.

- Decreased risk is most convincingly linked to fiber from vegetables, followed by fibers from non-soluble polysaccharides, starches, and fiber foods with carotenoids.

- Supplemental wheat bran in the amount of 13.5 grams per day can decrease the recurrence rate of adenomatous colon polyps.

- Since North Americans typically consume an average of only 8 to 15 grams of fiber per day, a dietary fiber supplement may be warranted.

- Currently, 40% of North Americans will develop cancer and the incidence is rising. If preventive measures are not instituted, the cost to the United States and its people will be enormous in terms of dollars, lost productivity, and lives lost.

#### AMOUNT OF FIBER

Depending on the study, North Americans typically consume only about 8-15 grams of fiber each day. Most of the consensus reports recommend 25 to 35 grams of fiber each day to protect against colorectal cancer. Unless North Americans have the time or inclination to become a grazing animal, it would be difficult to attain the protective level of fiber each day without taking a supplement.

#### CONCLUSION

- Dietary fiber is safe.
- Hundreds of studies, involving tens of thousands of subjects, demonstrate that 25 to 35 grams of dietary fiber daily can reduce the risk of colorectal cancer.

#### REFERENCES

1. Landis SH, Murray T, Bolden, Wingo PA: Cancer statistics, 1999. *CA - A Cancer J for Clinicians* 49(1):8-31, 1999.
2. Simone CB: *Cancer and Nutrition*. Garden City Park: Avery Publishing 1995.
3. Higginson J, Oettle AG: Cancer incidence in the Bantu and "Cape Colored" races of South Africa: report of a cancer survey in the Transvaal (1953-1955). *JNCI* 24:589-671, 1960.
4. Burkitt DP: Epidemiology of cancer of the colon and rectum. *Cancer* 28:3-13, 1971.
5. Alabaster O, Tang Z, Shivapurkar N: Dietary fiber and the chemopreventive modulation of colon carcinogenesis. *Mutat Res* 350(1):185-197, February 19, 1996.
6. Alberts DS, Lipkin M, Levin B: Genetic screening for colorectal cancer and intervention. *Int J Cancer* 69(1):62-63, February 20, 1996.
7. Alberts DS, Ritenbaugh C, Story JA, Aickin M, Rees-McGee S, Buller MK, Atwood J, Phelps J, Ramanujam PS, Bellapravalu S, Patel J, Bextinger L, Clark L: Randomized, double-blind, placebo-controlled study of effect of wheat bran fiber and calcium on fecal bile acids in patients with resected adenomatous colon polyps. *J Natl Cancer Inst* 88(2):81-92, January 17, 1996.

(cont. on p. 42)



8. Alberts DS, Einspahr J, Ritenbaugh C, Aickin M, Rees-McGee S, Atwood J, Emerson S, Mason-Liddil N, Bettinger L, Patel J, Bellapravalu S, Ramanujam PS, Phelps J, Clark L: The effect of wheat bran fiber and calcium supplementation on rectal mucosal proliferation rates in patients with resected adenomatous colorectal polyps. *Cancer Epidemiol Biomarkers Prev* 6(3):161-169, March 1997.
9. American Cancer Society: *Nutrition and Cancer: Cause and Prevention*. New York: American Cancer Society, 1984.
10. American Dietetic Association Position Paper: Health implications of dietary fiber. *J Am Diet Assoc* 97(10):1157-1159, October 1997.
11. Arbmán G, Axelsson O, Ericsson-Begodski AB, Fredriksson M, Nilsson E, Sjö Dahl R: Cereal fiber, calcium, and colorectal cancer. *Cancer* 15; 69(8):2042-2048, April 1992.
12. Barbare JC, Bedenne L, Bouche O, Cadiot G, Calais G, de Calan L, Conroy T, Ducreux M, Etienne PL, Elias D, Faivre J, Legoux JL, Maingon P, Mornex F, Nordlinger B, Ollier JC, Peiffert D, Pelletier G, Rat P, Rougier P, Ruskone-Fourmestral A, Seitz JF, Triboulet JP, Trinchet JC, Ychou M, *et al*: What can be done for digestive cancer in 1999: Recommendations of the French Foundation of Digestive Cancer (1st Part). *Gastroenterol Clin Biol* 23(4):502-511, April 1999.
13. Barbare JC, Bedenne L, Bouche O, Cadiot G, Calais G, de Calan L, Conroy T, Ducreux M, Etienne PL, Elias D, Faivre J, Legoux JL, Maingon P, Mornex F, Nordlinger B, Ollier JC, Peiffert D, Pelletier G, Rat P, Rougier P, Ruskone-Fourmestral A, Seitz JF, Triboulet JP, Trinchet JC, Ychou M, *et al*: What can be done for patients with digestive cancer in 1999: Guidelines of the French Foundation of Digestive Cancerology (2nd part). *Gastroenterol Clin Biol* 23(5):486-496, May 1999.
14. Biasco G: Chemoprevention of colorectal cancer. *Tumori* 83(1 Suppl):S5-S6, January/February 1997.
15. Bidoli E, Franceschi S, Talamini R, Barra S, La Vecchia C: Food consumption and cancer of the colon and rectum in north-eastern Italy. *Int J Cancer* 50(2):223-229, January 21, 1992.
16. Bingham SA: Epidemiology of dietary fibre and colorectal cancer: current status of the hypothesis. *Nutr Health* 4(1):17-23, 1985.
17. Bingham SA: Dietary fibre consumption and its association with large bowel cancer in man. *Med Oncol Tumor Pharmacother* 2(3):149-155, 1985.
18. Bingham SA: Non-starch polysaccharides as a protective factor in human large bowel cancer. *Princess Takamatsu Symp* 16:183-192, 1985.
19. Boutron-Ruault MC, Senesse P, Faivre J, Chatelain N, Belghiti C, Meance S: Foods as risk factors for colorectal cancer: a case-control study in Burgundy. *Eur J Cancer Prev* 8(3):229-235, July 1999.
20. Bright-See E, McKeown-Eyssen G, Jacobson EA, Newmark HW, Mathews R, Morson L, Jazmaji V, Bruce WR: Dietary fiber and cancer: a supplement for intervention studies. *Nutr Cancer* 7(4):211-220, 1985.
21. Braun EM: Diet factors and risk of the development of colorectal cancer. *Klin Khir* 6:44-46, 1996. Review. Russian.
22. Burkitt DP: Etiology and prevention of colorectal cancer. *Hospital Practice* 67-77, February 1984.
23. Calvert RJ, Klurfeld DM, Subramaniam S, Vahouny GV, Kritchevsky D: Reduction of colonic carcinogenesis by wheat bran independent of fecal bile acid concentration. *J Natl Cancer Inst* 79(4):875-880, October 1987.
24. Camire ME: Dietary fiber and colorectal cancer. *N Engl J Med* 340(3):169-176, January 21, 1999.
25. Caygill CP, Charlott A, Hill MJ: Relationship between the intake of high-fibre foods and energy and the risk of cancer of the large bowel and breast. *Eur J Cancer Prev* 7(Suppl 2):S11-S17, May 1998.
26. Chaplin MF: Bile acids, fibre and colon cancer: the story unfolds. *J R Soc Health* 118(1):53-61, February 1998.
27. Chlebowski RT: The scope of nutrition intervention trials with cancer-related endpoints. *Cancer* 74(9 Suppl):2734-2738, November 1, 1994.
28. ECP Consensus Panel on cereals and cancer: Consensus statement on cereals, fibre, and colorectal and breast cancers. Proceedings of the European Cancer Prevention Consensus meeting. Santa Margherita, Italy, October 2-5, 1997. *Eur J Cancer Prev* 7(Suppl 2):S1-S83, May 1998.
29. Cummings JH, Southgate DAT: Dietary fiber and colorectal cancer. *NEJM* 340(24):1925, 1999.
30. Cummings JH, Bingham SA, Heaton KW, Eastwood MA: Fecal weight, colon cancer risk, and dietary intake of nonstarch polysaccharides (dietary fiber). *Gastroenterology* 103(6):1783-1789, December 1992.
31. de Kok TM, van Faassen A, Bausch-Goldbohm R, ten Hoor F, Kleinjans JC: Fecapentaene excretion and fecal mutagenicity in relation to nutrient intake and fecal parameters in humans on omnivorous and vegetarian diets. *Cancer Lett* 62(1):11-21, February 14, 1992.
32. Earnest DL, Sampliner RE, Roe DJ, van Leeuwen B, Guillen J, Reid M, Martinez ME, Marshall JR, Alberts DS: Progress report: the Arizona phase III study of the effect of wheat bran fiber on recurrence of adenomatous colon polyps. *Am J Med* 106(1A):43S-45S, January 25, 1999.
33. Earnest DL, Einspahr JG, Alberts DS: Protective role of wheat bran fiber: data from marker trials. *Am J Med* 106(1A):32S-37S, January 25, 1999.
34. Edinboro PC, Lawrence W Jr, Nixon DW, Zfass A, Duckworth PF Jr, King R: Volunteers as adjunct researchers in a cancer prevention trial: the American Cancer Society Colon Polyp Prevention Study. *Cancer* 83(11):2384-2390, December 1, 1998.
35. European Organization for Cooperation in Cancer Prevention Studies, Provisional Dietary Recommendations: Joint ECP-IUNS Workshop and Third Annual Symposium on Diet and Human Carcinogenesis. *Nutr and Cancer*, 1986.
36. Evans DF: Physicochemical environment of the colon. *Eur J Cancer Prev* 7(suppl 2):S79-S80, 1998.
37. Faivre J, Bonithon-Kopp C: Chemoprevention of colorectal cancer. *Recent Results Cancer Res* 151:122-133, 1999.
38. Faivre J, Couillaud C, Kronborg O, Rath U, Giacosa A, De Oliveira H, Obrador T, O'Morain O: Chemoprevention of metachronous adenomas of the large bowel: design and interim results of a randomized trial of calcium and fibre. ECP Colon Group. *Eur J Cancer Prev* 6(2):132-138, April 1997.
39. Faivre J, Giacosa A: Primary prevention of colorectal cancer through fibre supplementation. *Eur J Cancer Prev* 7(Suppl 2):S29-S32, May 1998.
40. Faivre J: Colon cancer: epidemiology, pathological anatomy, diagnosis, course, treatment and prevention. *Rev Prat* 44(4):533-537, February 15, 1994.
41. Feldman EB: Dietary intervention and chemoprevention—1992 perspective. *Prev Med* 22(5):661-666, September 1993.
42. Food Guide Pyramid. Washington, D.C. U.S. Department of Agriculture, 1992. *Home and Garden Bulletin*, No 252.
43. Food, Nutrition, and the Prevention of Cancer: A Global Perspective. Washington, D.C.: American Institute for Cancer Research, 1998.
44. Ferguson LR, *et al*: Studies on the role of specific dietary fibres in protection against colorectal cancer. *Mutat Res* 350(1):173-184, February 19, 1996.
45. Franceschi S, Favero A, Parpinel M, Giacosa A, La Vecchia C: Italian study on colorectal cancer with emphasis on influence of cereals. *Eur J Cancer Prev* 7(Suppl 2):S19-23, May 1998.
46. Franceschi S, Parpinel M, La Vecchia C, Favero A, Talamini R, Negri E: Role of different types of vegetables and fruit in the prevention of cancer of the colon, rectum, and breast. *Epidemiology* 9(3):338-341, May 1998.
47. Freeman HJ: Role of high fibre foods in the prevention of colorectal neoplasia. *Can J Gastroenterol* 13(5):379-380, June 1999.
48. Freudenheim JL, Graham S, Marshall JR, Haughey BP, Wilkinson G: A case-control study of diet and rectal cancer in western New York. *Am J Epidemiol* 131(4):612-624, April 1990.
49. Fuchs CS, Giovannucci EL, Colditz GA, Hunter DJ, Stampfer MJ, Rosner B, Speizer FE, Willett WC: Dietary fiber and the risk of colorectal cancer and adenoma in women. *N Engl J Med* 340(3):169-176, January 21, 1999.
50. Garay CA, Engstrom PF: Chemoprevention of colorectal cancer: dietary and pharmacologic approaches. *Oncology (Huntingt)* 13(1):89-97; discussion 97-100, 105, January 1999.
51. Giovannucci E, Stampfer MJ, Colditz G, Rimm EB, Willett WC: Relationship of diet to risk of colorectal adenoma in men. *J Natl Cancer Inst* 84(2):91-98, January 15, 1992.
52. Glanz K, Kristal AR, Tilley BC, Hirst K: Psychosocial correlates of healthful diets among male auto workers. *Cancer Epidemiol Biomarkers Prev* 7(2):119-126, February 1998.
53. Graham S: Toward a dietary prevention of cancer. *Epidemiol Rev* 5:38-50, 1983.
54. Greenwald P: Colon cancer overview. *Cancer* 70(5 Suppl):1206-1215, September 1, 1992.
55. Greenwald P, Lanza E, Eddy GA: Dietary fiber in the reduction of colon cancer risk. *J Am Diet Assoc* 87(9):1178-1188, September 1987.
56. Handler S: Dietary fiber: can it prevent certain colonic diseases? *Postgrad Med* 73(2):301-307, February 1983.
57. Hardman WE, *et al*: Transforming growth factor alpha distribution in rectal crypts as a biomarker of decreased colon cancer risk in patients consuming cellulose. *Cancer Epidemiol Biomarkers Prev* 6(8):633-637, August 1997.
58. Harris PJ, Ferguson LR: Dietary fibres may protect or enhance carcinogenesis. *Mutat Res* 443(1-2):95-110, July 15, 1999.
59. Hassig CA, Tong JK, Schreiber SL: Fiber-derived butyrate and the prevention of colon cancer. *Chem Biol* 4(11):783-789, November 1997.
60. Heaton KW, Lewis SJ: Dietary fiber and colorectal cancer. *NEJM* 340(24):1925, 1999.
61. Heilbrun LK, Nomura A, Hankin JH, Stemmermann GN: Diet and colorectal cancer with special reference to fiber intake. *Int J Cancer* 44(1):1-6, July 15, 1989.
62. Held-Warmkessel J: Colon cancer: prevention and detection strategies. *Adv Nurse Pract* 6(7):42-45, July 1998.



63. Hill MJ: Cereals, cereal fibre and colorectal cancer risk: a review of the epidemiological literature. *Eur J Cancer Prev* 7(suppl 2): S5-S10, 1998.
64. Hill MJ: Cereals, cereal fibre and colorectal cancer risk: a review of the epidemiological literature. *Eur J Cancer Prev* 6(3):219-225, June 1997.
65. Hillman LC, Peters SG, Fisher CA, Pomare EW: Effects of the fibre components pectin, cellulose, and lignin on bile salt metabolism and biliary lipid composition in man. *Gut* 27(1):29-36, January 1986.
66. Howe GR, Benito E, Castelletto R, et al: Dietary intake of fiber and decreased risk of cancers of the colon and rectum: evidence from the combined analysis of 13 case-control studies. *JNCI* 84:1887-1896, 1992.
67. Hu JF, Liu YY, Yu YK, Zhao TZ, Liu SD, Wang QQ: Diet and cancer of the colon and rectum: a case-control study in China. *Int J Epidemiol* 20(2):362-367, June 1991.
68. Iscovich JM, L'Abbe KA, Castelletto R, Calzona A, Bernedo A, Chopita NA, Jmelnitzky AC, Kaldor J: Colon cancer in Argentina. I: Risk from intake of dietary items. *Int J Cancer* 51(6):851-857, July 30, 1992.
69. Ishikawa H: Interventional trial for cancer prevention. *Gan To Kagaku Ryobo* 26(2 Suppl):177-182, 1999.
70. Jansen MC, Bueno-de-Mesquita HB, Buzina R, Fidanza F, Menotti A, Blackburn H, Nissinen AM, Kok FJ, Kromhout D: Dietary fiber and plant foods in relation to colorectal cancer mortality: the Seven Countries Study. *Int J Cancer* 81(2):174, April 12, 1999.
71. Jeanteur P: Dietary fibre intake and colon cancer. *Bull Cancer* 86(7-8):611-613, July/August 1999.
72. Kahn MJ, et al: Chemoprevention for colorectal carcinoma. *Hematol Oncol Clin North Am* 11(4):779-794, August 1997.
73. Kenji T, Hedio O, Yasuharu O, Iwao Y, Tomohiro S, Katsuya Y, Kiichi M, Shigeru T, Hideki A: Dietary factors and prevention of colon cancer. *Nippon Geka Gakkai Zasshi* 99(6):368-372, June 1998.
74. Klipstein-Grobusch K, Kroke A, Voss S, Boeing H: Influence of lifestyle on the use of supplements in the Brandenburg nutrition and cancer study. *Z Ernahrungswiss* 37(1):38-46, March 1998.
75. Kritchevsky D: Cereal fibers and colorectal cancer: a search for mechanisms. *Eur J Can Prev* 7(suppl 2):S33-S39, 1998.
76. Kritchevsky D: Dietary fibre and cancer. *Eur J Cancer Prev* 6(5):435-441, October 1997.
77. Le Marchand L, Hankin JH, Wilkens LR, Kolonel LN, Englyst HN, Lyu LC: Dietary fiber and colorectal cancer risk. *Epidemiology* 8(6):658-665, November 1997.
78. Lipkin M, Reddy B, Newmark H, Lamprecht SA: Dietary factors in human colorectal cancer. *Annu Rev Nutr* 19:545-586, 1999.
79. Lubin F, Rozen P, Arieli B, Farbshtein M, Knaani Y, Bat L, Farbshtein H: Nutritional and lifestyle habits and water-fiber interaction in colorectal adenoma etiology. *Cancer Epidemiol Biomarkers Prev* 6(2):79-85, February 1997.
80. Lupton JR, Turner ND: Potential protective mechanism of wheat bran fiber. *Am J Med* 106(1A):24S-27S, 1999.
81. Macrae F: Wheat bran fiber and development of adenomatous polyps: evidence from randomized, controlled clinical trials. *Am J Med* 106(1A):38S-42S, January 25, 1999.
82. Madar Z, Stark A: Dietary fiber and colorectal cancer. *NEJM* 340(24):1926, 1999.
83. Martinez ME, Reid ME, Guillen-Rodriguez J, Marshall JR, Sampliner R, Aickin M, Ritenbaugh C, van Leeuwen B, Mason-Liddil N, Giuliano A, Vargas PA, Alberts DS: Design and baseline characteristics of study participants in the Wheat Bran Fiber trial. *Cancer Epidemiol Biomarkers Prev* 7(9):813-816, September 1998.
84. McKeigue PM, Adelstein AM, Marmot MG, Henly PJ, Owen RW, Hill MJ, Thompson MH: 85. Mohandas KM: Dietary fiber and colorectal cancer. *NEJM* 340(24):1925, 1999.
85. National Cancer Institute PDQ Screening/Prevention Summary for Health Professionals Prevention of Colorectal Cancer, January 1998, pp. 1-3.
86. National Cancer Institute PDQ Screening/Prevention Summary for Health Professionals Prevention of Colorectal Cancer, May 1998, pp. 1-14.
87. Negri E, Franceschi S, Parpinel M, La Vecchia C: Fiber intake and risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev* 7(8):667-671, August 1998.
88. Nelson RL, et al: Determination of factors responsible for the declining incidence of colorectal cancer. *Dis Colon Rectum* 42(6):741-752, June 1999.
89. Nordgaard I, et al: Colonic production of butyrate in patients with previous colonic cancer during long-term treatment with dietary fibre (Plantago ovata seeds). *Scand J Gastroenterol* 31(10):1011-1020, October 1996.
90. Otto C, et al: Nutrition and cancer: nutrition-associated risk for the development of various malignancies. *Fortschr Med* 113(17):267-271, June 20, 1995.
91. Owen RW, Spiegelhalter B, Bartsch H: Phytate, reactive oxygen species and colorectal cancer. *Eur J Can Prev* 7(suppl 2):S41-S54, 1998.
92. Potter JD: Nutrition and colorectal cancer. *Cancer Causes Control* 7(1):127-146, January 1996.
93. Primary prevention of colorectal cancer and polyps: does fiber have a role? Proceedings of a symposium. New York City, New York, December 2, 1997. *Am J Med* 106(1A):1S-51S, January 5, 1999.
94. Rafter J, Glinghamer B: Interaction between the environment and genes in the colon. *Eur J Can Prev* 7(suppl 2):S69-S74, 1998.
95. Ravin ND: Dietary fiber and colorectal cancer. *NEJM* 340(24):1924-1925, 1999.
96. Reddy BS: Role of dietary fiber in colon cancer: an overview. *Am J Med* 106(1A):16S-19S, discussion 50S-51S, January 25, 1999.
97. Sandler RS: Epidemiology and risk factors for colorectal cancer. *Gastroenterol Clin North Am* 25(4):717-735, December 1996.
98. Schatzkin A, et al: The polyp prevention trial I: rationale, design, recruitment, and baseline participant characteristics. *Cancer Epidemiol Biomarkers Prev* 5(5):375-383, May 1996.
99. Scheppach W, Bingham S, Boutron-Ruault MC, Gerhardtsson de Verdier M, Moreno V, Nagengast FM, Reifen R, Riboli E, Seitz HK, Wahrendorf J: WHO consensus statement on the role of nutrition in colorectal cancer. *Eur J Cancer Prev* 8(1):57-62, February 1999.
100. Slavin JL: Implementation of dietary modifications. *Am J Med* 106(1A):46S-49S, 1999.
101. Smith JG, Yokoyama WH, German JB: Butyric acid from the diet: actions at the level of gene expression. *Crit Rev Food Sci Nutr* 38(4):259-297, May 1998.
102. Statland BE: Nutrition and cancer. *Clin Chem* 38(8 Pt 2):1587-1594, August 1992.
103. Steinmetz KA, Potter JD: Vegetables, fruit, and cancer. I. Epidemiology. *Cancer Causes Control* 2(5):325-357, September 1991.
104. Szilagyi A: Altered colonic environment, a possible predisposition to colorectal cancer and colonic inflammatory bowel disease: rationale of dietary manipulation with emphasis on disaccharides. *Can J Gastroenterol* 12(2):133-146, March 1998.
105. Thun MJ, Calle EE, Namboodiri MM, Flanders WD, Coates RJ, Byers T, Boffetta P, Garfinkel L, Heath CW Jr: Risk factors for fatal colon cancer in a large prospective study. *J Natl Cancer Inst* 84(19):1491-1500, October 7, 1992.
106. Trock B, Lanza E, Greenwald P: Dietary fiber, vegetables, and colon cancer: critical review and meta-analyses of the epidemiologic evidence. *J Natl Cancer Inst* 82(8):650-661, April 18, 1990.
107. Weisburger JH: Worldwide prevention of cancer and other chronic diseases based on knowledge of mechanisms. *Mutat Res* 402(1-2):331-337, June 18, 1998.
108. Zimbalist EH, et al: Genetic and environmental factors in colorectal carcinogenesis. *Dig Dis* 13(6):365-375, November/December 1995.



**Charles B. Simone, M.D.**, is director and founder of the Simone Protective Cancer Center in Lawrenceville, NJ. Dr. Simone is an internist, medical oncologist, tumor immunologist, and radiation oncologist. Dr. Simone wrote *Cancer and Nutrition: A Ten Point Plan to Reduce Your Risk of Getting Cancer* (McGraw-Hill, 1981; revised Avery, 1995). He also wrote *Breast Health* (Avery, 1995), and *Shark Cartilage and Cancer* (Paradigm Books, 1995). He can be reached at mail@DrSimone.com, and his website is www.DrSimone.com.



**Nicole L. Simone, B.S.E.**, graduated from the bioengineering department of the University of Pennsylvania. She is currently a medical student at UMDNJ, and is the co-manager of the Laser Capture Microdissection (LCM) Lab. She is also co-director of the Proteomics Branch of the Cancer Genome Anatomy Project at the National Cancer Institute, National Institutes of Health. She has written several articles for journals such as *Science* and *The Lancet*. In addition, she is co-inventor of a technique for protein analysis from microdissected tissue.



**Charles B. Simone, II**, attends the University of Pennsylvania. He has written articles for journals such as *The Lancet*. He is the recipient of numerous awards, including the *USA Today* National High School Academic All-Star Team, and Toyota Community Scholars Regional Scholarship. Nicole and Chuck help educate cancer patients and their families about healthy lifestyle modification. They both helped to spearhead the KID START program, designed to teach children, through seminars and health fairs, about healthful lifestyles.